

# Calf Navel Ill

## SOP

*This is a farm specific Standard Operating Procedure (SOP) drawn up by Firmount Veterinary Clinic for the Cafre dairy centre Herd. Farmers should consult with their veterinary practitioner to develop SOP's appropriate to their herd.*

Inject 1ml *Noroclav* per 25kg body weight into the muscle once daily for 5 days. On 1st 3rd and 5th days of treatment also give an injection of 2.5ml *loxicom* per 100kg under the skin and record all treatments in medicines book.

After 5 days of treatment  
have signs of navel ill gone?

Yes

No action  
required

No

Consult  
Vet

**Withdrawal Periods:**

Noroclav: meat- 42 days

Loxicom: meat- 15 days

# Calf joint ill

## SOP

*This is a farm specific Standard Operating Procedure (SOP) drawn up by Firmount Veterinary Clinic for the Cafre Dairy Centre Herd. Farmers should consult with their veterinary practitioner to develop SOP's appropriate to their herd.*

Inject 1ml *Noroclav* per 25kg into the muscle once daily for 5 days. On 1<sup>st</sup> 3<sup>rd</sup> and 5<sup>th</sup> days of treatment also give an injection of 2.5ml *Loxicom* per 100kg under the skin and record all treatments in medicines book.

After 5 days of treatment  
have signs of joint ill gone?

No

No action  
required

Yes

Consult  
Vet


**Withdrawal Periods:**

**Noroclav: meat-42 days**

**Loxicom: meat-15 days**

# Calf Scour SOP

This is a farm specific Standard Operating Procedure (SOP) drawn up by Firmount Veterinary Clinic for the Cairn Dairy Centre Herd. Farmers should consult with their veterinary practitioner to develop SOP's appropriate to their herd.



**Withdrawal Periods:**

Glutalyte: 0 days

# Pneumonia in young cattle SOP

This is a farm specific Standard Operating Procedure (SOP) drawn up by Firmount Veterinary Clinic for the Cafre Dairy Centre Herd. Farmers should consult with their veterinary practitioner to develop SOP's appropriate to their herd.



**Withdrawal Periods:**

Hexasol LA      meat-35days